

Comments on evaluation of ²⁶Al Electron-Capture and Positron Decay Data by Shiu-Chin Wu and E. Browne

The *Limitation of Relative Statistical Weight* [2] (LWM) method, used for averaging numbers throughout this evaluation, provided a uniform approach for the analysis of discrepant data. The uncertainty assigned to the recommended values was always greater than or equal to the smallest uncertainty of the values used to calculate the average.

Decay Scheme

²⁶Al decays 100% by EC + β^+ to ²⁶Mg 2+ states at 1808.72 and 2938.41 keV. A measured relative emission probability for the positron annihilation radiation (511 keV) produced a total β^+ branching of 82(2)% (72Sa02). This branching agrees well with a value of 81.7% predicted by theory[1].

Nuclear Data

The recommended half-life of ²⁶Al, $7.17(24) \times 10^5$ y, is a weighted average ($\chi^2/N-1=0.64$, LWM) of $7.16(32) \times 10^5$ y, from partial $T_{1/2}(\beta^+)=8.73(30) \times 10^5$ y, β^+ counting and mass spectrometric analysis [3], combined with $\beta^+(\%)=82(2)$ from measurement of positron annihilation radiation (72Sa02)); $7.05(24) \times 10^5$ y, by specific activity and mass spectrometric analysis [4]; $7.8(5) \times 10^5$ y, by specific activity from a source produced by ²⁶Mg(p,n) ²⁶Al (the number of ²⁶Al atoms in the source were estimated by using the reaction cross section) [5]; and $7.02(56) \times 10^5$ y, by counting the atoms of ²⁶Al that did not disintegrate in the source (Accelerator Mass Spectrometry) [6]. The internal uncertainty in the weighted average is 0.17×10^5 y. However, we adopted a conservative value of 0.24×10^5 y. (the lowest uncertainty in the input experimental half-life results), because of possible correlations between the measured values.

Gamma Rays

Gamma-ray energies and relative emission probabilities are from measurements with Ge(Li) detectors (72Sa02). Since EC + β^+ feeding to the ground state of ²⁶Mg is not expected ($\Delta J=5$), the sum of the relative emission probabilities of the 1808.65 (0.9976) and 2938 keV (0.025(2)) gamma rays to the ground state of ²⁶Mg was used to normalize the decay scheme (See Table 1). A conservative estimate based on data reported by 73Ra10 gives $\text{Log } ft > 24$ for a fourth forbidden unique transition. This value corresponds to $I_{EC} < 0.0005\%$ for a possible (but yet unobserved) EC transition to ²⁶Mg ground state. In this calculation we assumed a fractional uncertainty of 2% for the relative emission probability of the 1808.65 keV gamma ray. This value, which is based on the fractional uncertainty of the annihilation radiation (1.641(32)) quoted by 72Sa02, has a negligible effect on the uncertainty of its absolute gamma-ray emission probability.

Table 1 : Gamma-ray Energies and Relative Emission Probabilities from ²⁶Al EC + β^+ Decay

Energy (keV)	Relative Emission Probability (per disint.)
annihil. rad.	1.641(32) *
1129.67(10)	0.025(2)
1808.65(7)	0.9976
2938	0.0024(4)

* Corrected for annihilation of positrons in flight

Gamma-ray multipolarities and mixing ratios are from 78En08. Conversion coefficients, are insignificant, however, theoretical interpolated values between Z=10 and Z=14 from 76Ba63 have been included in this evaluation.

Electron Capture and Positron Emission (β^+)

Electron-capture and β^+ end-point energies given in Tables 2.2 and 2.3 are equal to $Q_{EC} = 4004.19(6)$ (95Au04) minus the individual level energies, and to the electron-capture energies minus $2 m_0c^2$ (1022 keV), respectively. EC + β^+ feedings to the 1808.72 keV and 2938.41 keV levels are from gamma-ray emission probability balances. The individual electron-capture and positron emission probabilities are based on theoretical [1] β^+ / EC ratios. The total measured positron absolute emission probability (82(2)%) agrees well with a value of 81.7%, calculated for a second forbidden unique transition using a theoretical $\beta^+ / (EC + \beta^+)$ ratio of 0.840. β^+ (P_{β^+}) and EC (P_{EC}) are given in percent (%) on the decay scheme. Fractional atomic shell electron-capture probabilities are theoretical values [7], calculated with the LOGFT computer program [8].

Atomic Data

The X-ray and Auger electron emission probabilities given in section 3 are values calculated by using the computer program RADLST [9], the electron capture probabilities from section 2.2, and atomic data from 96Sc06.

Acknowledgments

The authors are grateful to E.B. Norman and R.W. Kavanagh for enlightening discussions regarding the half-life of ²⁶Al.

References

- [1] - P.F.Zweifel, *Capture-Positron Branching Ratios*, Phys. Rev. **96**, 1572 (1954); P.F.Zweifel, *Allowed Capture-Positron Branching Ratios*, Phys. Rev. **107**, 329 (1957)
- [2] - M.J. Woods and A.S. Munster, *Evaluation of Half-Life Data*. National Physical Laboratory, Teddington, UK, Rep. RS(EXT) 95, (1988).
- [3] - R.A. Rightmire, J.R. Simanton, and T.P. Kohman, *Über die Halbwertszeit des langlebigen ²⁶Al*, Phys. Rev. **113**, 1069 (1959)
- [4] - T.L. Norris, A.J. Gancarz, D.J. Rokop and K.W. Thomas, *Half-Life of ²⁶Al*, Proc. of the Fourteenth Lunar and Planetary Science Conf., J. Geophysical Research, **88**, B331 (1983)
- [5] - J.H. Thomas, R.L. Rau, R.T. Skelton and R.W. Kavanagh, *Half-life of ²⁶Al*, Phys. Rev. **C30**, 385 (1983)
- [6] - R. Middleton, J. Klein, G.M. Raisbeck and F. Yiou, *Accelerator Mass Spectrometry with ²⁶Al*, Nucl. Inst. and Meth., **218**, 430 (1983)
- [7] - W. Bambynek, H. Behrens, M.H.Chen, B. Crasemann, M.L.Fitzpatrick, K.W.D. Ledingham, H. Genz, M. Mutterer and R. L. Intemann, *Orbital Electron Capture by the Nucleus*, Rev. Mod. Phys. **49**, 77 (1977)
- [8] - *The Program LOGFT*, National Nuclear Data Center, September 3, 1993.
- [9] - *The Program RADLST*, Thomas W. Burrows, report BNL-NCS-52142, February 29, 1988.
- 72Sa02 - E.A. Samworth, E.K. Warburton, G.A.P. Engelbertink, *Beta Decay of the ²⁶Al Ground State*. Phys. Rev. **C5**, 138 (1972).
- 76Ba63 - I.M. Band, M.B. Trzhaskovskaya, M.A. Listengarten, *Internal Conversion Coefficients for Atomic Numbers Z < 30*, At. Data Nucl. Data Tables **18**, 433 (1976).
- 78En08 - P.M. Endt and C. van der Leun, *Energy Levels of A = 26*, Nucl. Phys. **A310**,156 (1978).
- 90En08 - P.M. Endt, *Energy Levels of A = 26*, Nucl. Phys. **A521**, 155 (1990).
- 95Au04 - G. Audi and A.H. Wapstra, *The 1995 Update to the Atomic Mass Evaluation*. Nucl. Phys. **A595**, 409 (1995).
- 96Sc06 - E. Schönfeld and H. Janßen, *Evaluation of Atomic Shell Data*, Nucl. Instr. Meth. in Phys. Res. **A369**, 527 (1996).