

²¹⁸At - Comments on evaluation of decay data by V. Chisté and M. M. Bé

This evaluation was completed in 2007. Literature available by January 2007 was included.

1 Decay Scheme

²¹⁸At disintegrates by alpha emission (99.9 (1) %) to ²¹⁴Bi mainly. The γ transitions between the ²¹⁴Bi levels have not been observed. A weak beta minus emission (0.1 (1) %) to Rn-218 has been pointed out (1948Wa20). Spins and parities are from the mass-chain evaluation of Y. A. Akovali (1987El12, 1995El08 for A = 218 and 1995El07 for A = 214) and A. K. Jain (2006Ja03 for A = 218).

2 Nuclear Data

The Q values (α and β^-) are from the atomic mass evaluation of Audi *et al.* (2003Au03).

Experimental ²¹⁸At half-life values (in seconds) are given in Table 1:

Table 1: Experimental values of ²¹⁸At half-life.

Reference	Experimental value (s)	Comments
R. J. Walen (1949Wa05)	1.3 (2)	Uncertainty increased to take into account systematic uncertainty.
D. G. Burke (1989Bu09)	1.5 (3)	
Recommended value	1.4 (2)	$\chi^2 = 0.31$

The original uncertainty value given by R. J. Walen (1949Wa05) was multiplied by 2, in order to take into account the systematic uncertainties which were not considered by 1949Wa05. A weighted average has been calculated using Lweight computer program (version 3).

The recommended value of ²¹⁸At half-life is the weighted average of **1.4** seconds with an internal uncertainty of 0.2 seconds. The reduced- χ^2 value is 0.31.

2.1 a Transitions and Emissions

The energies of the α -particle transitions given in Section 2.1 were calculated from Q_α (2003Au03) and level energies.

The energy of $\alpha_{0,0}$, $\alpha_{0,1}$ and $\alpha_{0,2}$ emissions given in section 3 were measured by R.J. Walen (1963Wa29 (see 1964Hy02) and 1958Wa16), the adopted values are those recommended by A. Rytz (1991Ry01) where the original energy was decreased by 1 keV, due to a change in calibration energy (1995El07).

The $\alpha_{0,0}$, $\alpha_{0,1}$ and $\alpha_{0,2}$ emission probabilities are the measured values of R. J. Walen (1958Wa16), 3.6, 90.0 and 6.4 respectively, without uncertainties. From R. J. Walen, the total α decay is 99.9 (1) %. Since, there is no precision in the Walen's paper, the uncertainty of 0.1 % from propagation of the β^- transition probability (1948Wa20) has been assigned to each α line.

2.2 β^- Transitions and Emissions

The maximum energy of the β^- transition in the decay of $^{218}\text{At} \rightarrow ^{218}\text{Rn}$ is given by Audi (2003Au03) and, without any other information available, is affected to a ground state to ground state transition.

The adopted β^- transition probability was measured by R. J. Walen (1948Wa20) to be 0.1 (1) %

3 References

- 1948Wa20 – R. J. Walen, *Comp. Rend. Acad. Sci. (Paris)* 227(1948)1090 [Branching ratio].
1949Wa05 – R. J. Walen, *J. Phys. Radium* 10(1949)95 [I_β , Half-life].
1958Wa16 – R. J. Walen, G. Bastin, *Comp. Rend. Int. Conf. Nucl. Phys., Paris, (1959)*910 [E_α , I_α , I_β].
1963Wa29 – R. J. Walen, G. Bastin-Scoffier, *Priv. Comm. quoted by 1964Hy02 (1963)* [I_α].
1964Hy02 – E. K. Hyde, I. Perlman, G. T. Seaborg, *The nuclear properties of heavy elements, Vol. II*, Prentice-Hall, Inc., Englewood Cliffs, N.J., (1964)460 [I_α].
1979Ry03 – A. Rytz, *At. Data and Nucl. Data Tables* 23(1979)205 [E_α , I_α].
1987El12 – Y. A. Ellis-Akovali, *Nucl. Data Sheets* 52(1987)789 [I_α , E_α , spin and parity].
1989Bu09 – D. G. Burke, H. Folger, H. Gabelmann, E. Hagebø, P. Hill, P. Hoff, O. Jonsson, N. Kaffrell, W. Kurcewicz, G. Løvholden, K. Nybø, G. Nyman, H. Ravn, K. Riisager, J. Rogowski, K. Steffensen, T. F. Thorsteinsen and ISOLDE Collaboration, *Z. Phys.* A333(1989)131 [Half-life].
1991Ry01 – A. Rytz, *At. Data and Nucl. Data Tables* 47(1991)205 [E_α , I_α].
1995El07 – Y. A. Akovali, *Nucl. Data Sheets* 76(1995)127 [I_α , E_α , spin and parity].
1995El08 – Y. A. Akovali, *Nucl. Data Sheets* 76(1995)457 [I_α , E_α , spin and parity].
2003Au03 – G. Audi, A. H. Wapstra, C. Thibault, *Nucl. Phys.* A729(2003)129 [Q].
2006Ja03 – A. K. Jain, B. Singh, *Nucl. Data Sheets* 107(2006)1027 [I_α , E_α , spin and parity].