

¹²⁹I - Comments on evaluation of decay data by V. P. Chechev and V. O. Sergeev

1- Decay Scheme

The 2nd unique forbidden β^- -transition to the 1/2⁺ ground state of ¹²⁹Xe was not observed. In 1954 Der Matiosian and Wu (1954De17) showed experimentally that this β^- -branch intensity did not exceed 1 %. This limit gives a $\log f_{2ut} = 14.9$ (or $\log f_{0t} = 15.8$), which is consistent with the $\log f_{2ut}$ values of 14.6 – 15.8 tallied in 1998Si17 for ten cases from A=22 to A=138, excluding ¹⁰Be, with 13.8, and ²⁰⁹Po, with 14.36. The highest value of 15.8 corresponds to 0.13% for the transition considered.

Therefore, we have adopted the probability of the 2nd unique forbidden β^- -transition to the 1/2⁺ ground state of ¹²⁹Xe $P(\beta^-_{0,0}) = 0.05(5)\%$ with the uncertainty which provides the limits from 0 to 1% according to 1954De17.

2- Nuclear Data

The Q value has been computed on the basis of the spectrometric measurement of the $\beta^-_{0,1}$ energy by N. Coursol (1979CoZG) and the evaluated gamma-ray energy. This measurement gives a more accurate Q value than 194(3) keV, presented in the atomic mass evaluation (1995Au04).

The following four experimental values for the ¹²⁹I half-life are available (in units of 10⁷ years).

1.72(9)	1951 Ka16
1.56(6)	1957Ru65
1.57(4)	1972Em01
1.97(14)	1973Ku17

Use of the LRSW method leads to a higher uncertainty (0.047) in 1972Em01. Our recommended value has been obtained as the weighted mean with the external uncertainty 0.06 expanded due to the Student's factor (or MBAYS uncertainty) : 1.61(7). Thus our recommended value for the ¹²⁹I half-life is 1.61(7) × 10⁷ years.

2.1. β^- -Transitions

The energy of the $\beta^-_{0,1}$ transition has been adopted from 1979CoZG (Coursol). For the probabilities $P(\beta^-_{0,1})$ and $P(\beta^-_{0,0})$ see discussion in sect.1.Decay Scheme.

2.2. Gamma-ray Transitions and Internal Conversion Coefficients

The correction for recoil has not changed the γ -ray transition energy.

The emission probability of the γ -ray transition (photons + electrons) has been adopted as 99.5(5)%.(see discussion in sect.1).

The multipolarity of the γ -ray transition was measured in 1965Ge04 (M1) and 1974Ra26 (M1 + 0.073(27)% E2).

ICC's have been interpolated from theoretical values of 1978Ro22 for the adopted multipolarity of $M1 + 0.07(3)\% E2$. The uncertainties in the theoretical values are as follows: 1% for α_K and 3% for α_L , α_M , α_{NO} . The ratio α_{NO}/α_M has been taken from 1971Dr11. The ICC interpolated from other tables (1968Ha53, 1969Ha61, 1978Band) agree with the adopted values within the limits of the stated uncertainties.

The interpolated value $\alpha_K^{\text{theory}} = 10.59(11)$ can be compared with the following experimental values: 10.6 (1968ReZY), 9.8(9) (1970Gy01), 10.2(4) (1970SaZI), 10.2(5) (1977Ra23), and 10.6(4) (1985Ba73), which have an unweighted average of 10.3.

3. Atomic Data

3.1. Fluorescence yields

The fluorescence yields have been taken from 1996Sc06 (Schönfeld and Janßen).

3.2. X rays

X-ray energies are based on the wavelengths given in the compilation of 1967Be65 (Bearden).

The relative K x-ray emission probabilities have been taken from 1996Sc06 and 1999Schönfeld.

3.3. Auger Electrons

The energies of Auger electrons are from 1977La19 (Larkins) and 1998Schönfeld.

The ratios $P(KLX)/P(KLL)$ and $P(KXY)/P(KLL)$ have been taken from 1996Sc06.

4. Electron emissions

The energies of the conversion electrons have been calculated from the γ -ray transition energy given in sect. 2.2 and the electron binding energies. Their absolute emission probabilities have been calculated using the conversion coefficients given in 2.2 and the absolute γ -ray emission probability.

For the L-shell the ratios $L_1:L_2:L_3 = 100:8.9(4):3.13(14)$ obtained from theoretical conversion coefficients can be compared with the experimental $L_1:L_2:L_3 = 100:10.0(4):3.1(3)$ from $^{129}\text{Cs} \rightarrow ^{129}\text{Xe}$ decay (1965Ge04).

Values of the emission probabilities of K-Auger electrons have been calculated using our recommended $P(\text{ceK})$ and $P(\text{ceL})$ values and atomic data given in 3.1.

The maximum energy of β^- particles with energy of 151 keV has been taken from 1979CoZG(Coursol). The average energy of β^- particles calculated with the LOGFT program, which uses an allowed spectral shape, is 40.6(3) keV. The SPEBETA program gives a different value of 37 keV (2001 Be). In 2001Be the shape factor $C(W) = q^2 + (0.10 \pm 0.01)p^2$ was used that given by E. der Matiosian and C. S. Wu (1953DE10) (measurement with a magnetic spectrometer). The value of 37 keV is supported also by the calculation of Kolobachkin et.al. (See the book "Beta emissions of fission products", authors: V. M. Kolobachkin, P. M. Rubtsov, V. G. Alexankin and P. A. Ruzhanskiy . – Moscow, Atomizdat, 1978, p.189. In Russian). They found 36 keV for the average energy of β^- particles of ^{129}I . So we adopt 37(1) keV as the recommended value.

5. Photon Emissions

5.1 X-Ray Emissions

Our recommended value for the total K x-ray absolute emission intensity has been calculated as $P_{\text{XK}}^{\text{eval.}} = \omega_{\text{K}}\alpha_{\text{K}}P_{\gamma}(39.6) = 69.8(11)\%$, based on the adopted value of ω_{K} , a theoretical value of α_{K} , and our recommended value of $P_{\gamma}(39.6) = 7.42(8)\%$. This K x-ray emission probability agrees well with the result of the measurement $P_{\text{XK}}^{\text{exp.}} = 70.2(8)\%$ in 1985Ba73, relative to $P_{\gamma}(39.6) = 7.46\%$ (or $69.8(8)\%$, relative to $P_{\gamma}(39.6) = 7.42\%$), and it also agrees with the less accurate experimental result from 1977Ra23: $73(6)\%$.

The absolute emission probabilities of the K x-ray components have been deduced from the total P_{XK} using the relative probabilities from sect. 3.2.

The total absolute emission probability of L x-rays has been deduced using the adopted values of $\overline{\omega}_{\text{L}}$ and n_{KL} and the recommended values of $P(\text{ce}_{\text{K}}) = 78.6(12)$ and $P(\text{ce}_{\text{L}}) = 10.8(4)\%$.

5.2. Gamma Emissions

A γ -ray energy of 39.578(4) keV has been adopted from 1985Ba73 from an accurate measurement made with a planar HPGe detector. The adopted value coincides with 39.578(2) keV for the energy of the first excited level in ^{129}Xe (1996Te01), deduced from the decay of ^{129}Cs .

Other less accurate experimental values of $E(\gamma_{1,0})$ are (in keV): 39.58(3) (1965Ge04), 39.6(2) (1966Re10), 39.4(3) (1967Gr05), 39.58(5) (1972Ta15), and 39.581(15) (1976Me16).

The absolute γ -ray emission probability (P_{γ}) has been computed as $P(\beta_{1,0})/(1+\alpha_{\text{T}})$. The uncertainty in P_{γ} includes the uncertainty of 0.5% in $P(\beta_{1,0})$, and 1% in α_{T} .

References

- 1951KA16 S. Katcoff, O. A. Schaeffer, J. M. Hastings, Phys. Rev. 82(1951)688. (Half-life)
 1953DE10 Der E. Mateosian, C. S. Wu, Phys. Rev. 91(1953)497A. (Beta spectrum shape factor)
 1954DE17 Der E. Mateosian, C. S. Wu, Phys. Rev. 95(1954)458. (Gamma ray energy)
 1957RU65 H. T. Russell, Report ORNL – 2293(1965). (Half-life)
 1965GE04 J. S. Geiger, R. L. Graham, I. Bergstrom, F. Brown, Nucl. Phys. 68(1965)352. (Gamma ray energy, multipolarity)
 1966RE10 I. Rezanka, A. Spalek, J. Frana, A. Mastalka, Nucl. Phys. 89(1966)609. (Gamma ray energy)
 1967BE65 J. A. Bearden, Revs. Modern Phys. 39(1967)78. (X-ray energies)
 1967GR05 G. Graeffe, W. B. Walters, Phys. Rev. 153(1967)1321. (Gamma ray energy, K ICC)
 1968HA53 R. S. Hager, E. C. Seltzer. Nucl. Data Tables A4(1968)1. (Theoretical ICC)
 1968REZY S. A. Reynolds, J. F. Emery, ORNL-4343 (1968) p 78. (K ICC)
 1969HA61 R. S. Hager, E. C. Seltzer, Nucl. Data Tables A6(1969)1. (Theoretical ICC)
 1970GY01 F. N. Gyax, R. F. Jenefsky, H. J. Leisi, Phys. Letters 32B(1970)359. (K ICC)
 1970SAZI K. S. R. Sastry, R. E. Wood, J. M. Palms, P. V. Rao, Bull. Am. Phys. Soc. 15(4), JE12(1970)623. (K ICC)
 1971DR11 O. Dragoun, Z. Plajner, F. Schmutzler, Nucl. Data Tables A9(1971)119. (Theoretical ICC)
 1972Em01 J. F. Emery, S. A. Reynolds, E. I. Wyatt, G. I. Gleason, Nucl. Sci. Eng. 48(1972)319. (Half-life)

- 1972TA15 H. W. Taylor, B. Singh, J. Phys. Soc. Japan 32(1972)1472. (Gamma ray energy)
- 1973KU17 J. G. Kuhry, G. Bontems, Radiochem. Radioanal. Letters 15(1973)29. (Half-life)
- 1974MA24 G. Marest, R. Haroutunian, I. Berkes, M. Meyer, M. Rots, J. De Raedt, H. Van de Voorde, Van de, H. Oonis, R. Coussement, Phys. Rev. C10(1974)402. (Multipolarity)
- 1974RA26 K. Venkata Ramaniah, T. Seshi Reddy, K. Venkata Reddy, Current Sci. (India) 43 (1974)406. (Multipolarity, K ICC)
- 1976ME16 R. A. Meyer, F. F. Momyer, J. H. Landrum, E. A. Henry, R. P. Yaffe, W. B. Walters, Phys. Rev. C14(1976)1152. (Gamma ray energy)
- 1977RA23 T. K. Ragimov, D. F. Rau, V. I. Timoshin, Bull. Akad. Sci. USSR, Phys. Ser. (1977), 1941(6), 97. Izv Akad Nauk SSSR. Ser. Fiz. 41(1977)1222 (K ICC)
- 1979COZG N. F. Coursol, Thesis, Univ de Paris (1979). (K ICC, M ICC, β^- emission energy)
- 1985BA73 G. Barci-Funel, M. C. Kouassi, G. Ardisson, Nuclear Instrum. Methods 24(1985)252 (K ICC, gamma ray energy, gamma ray and K X-ray emission probabilities)
- 1995AU04 G. Audi and A. H. Wapstra, Nucl.Phys. A595(1995)409. (Q value)
- 1996SC06 E. Schönfeld, H. Janßen, Nucl. Instrum. Methods Phys. Res. A369(1996)527. (Atomic data)
- 1996TE01 Y. Tendow, Nucl. Data Sheets 77(1996)631 (Level energies)
- 1998Schönfeld E. Schönfeld and G. Rodloff, report PTB-6,11-98-1, Braunschweig, October 1998. (Energies of Auger electrons)
- 1998Si17 B. Singh, J. L. Rodriguez, S. S. M. Wong, J. K. Tuli, Nucl. Data Sheets 84(1998)487. (Systematics of log ft values)
- 1999Schönfeld E. Schönfeld and G. Rodloff, report PTB-6.11-1999-1999-1, Braunschweig, February 1999. (KX ray energies and relative emission probabilities)
- 2001Be M. M. Bé, INDC(NDS)-422, IAEA, Vienna, P.112. (Average β^- energy)